The Marine Resources of the Parker River-Plum Island Sound Estuary: An Update after 30 Years

edited and compiled by

Robert Buchsbaum, Massachusetts Audubon Society

Tim Purinton, Parker River Clean Water Association and
Massachusetts Audubon Society
and
Britta Magnuson, Massachusetts Audubon Society

funded by the Massachusetts Office of Coastal Zone Management

Based on the 1968 monograph by the Massachusetts Division of Marine Fisheries:

A Study of the Marine Resources of the Parker River-Plum Island Sound Estuary by

William Jerome, Arthur Chesmore, and Charles O. Anderson, Jr.

and on the

Plum Island Sound Minibay Project Report

by

Robert Buchsbaum, Andrea Cooper, and Joan LeBlanc

TABLE OF CONTENTS

CHAPTER 1. INTRODUCTION	1
Personnel and Acknowledgements	1
1.1. Objectives	3
1.2. Study Area	3
CHAPTER 2. GEOLOGY AND MORPHOLOGY OF THE PARKER RIVER-PLUM	[
ISLAND SOUND ESTUARY	5
2.1. Overview	
2.2. Methods	
2.3. Geography and Morphometry	
2.4. Geological Background	9
CHAPTER 3. THE WATER QUALITY AND FLUSHING CHARACTERISTICS OF	7
THE PARKER RIVER-PLUM ISLAND SOUND ESTUARY	
3.1. Overview	13
3.2. Methods and Materials	13
3.3. Results and Discussion of Water Quality Analysis	
3.4. Toxic Contaminants in the Plum Island Sound Region	
3.5. Pollution and Its Effect on Marine Resources – A Thirty Year Perspective	
CHAPTER 4. FISH	75
4.1. Changes in the fish community of Plum Island Sound from 1965 through 1994	75
	103
4.3. Striped Bass	108
4.4. Smelt	110
4.5. White Perch	111
	111
4.7. Summary	113
CHAPTER 5. SHELLFISH	115
	115
5.2. Surf Clams	132
5.3. Razor Clams	132
	132
•	133
5.6. Sea Worms, Family Nereidae and Family Glyceridea	133
5.7. American Lobster	
134	
5.8. Green Crab	134
5.9. Quahogs	134
5.10. Summary	134
CHATPER 6. BIRDS AT PLUM ISLAND: A COMPARISON OF PRESENT AND	
HISTORICAL OBSERVATIONS	136
6.1. Background	
6.2. Methods	
6.3. Results	138
CHAPTER 7. THE VEGETATION OF PLUM ISLAND	143
7.1. Types of plants occurring in the Plum Island Sound region	

7.2. The Salt Hay Industry		
CHAPTER 8. HABITAT ISSUES FACED BY THE PLUM ISLAND SOUND		
ESTUARY	148	
8.1. Water Quality Decline and Eutrophication	148	
8.2. Impacts of Tidal Restrictions on Vegetation		
8.3. Salt Marsh Mosquito Control		
8.4. Restoration of Anadromous Fish Habitat	155	
8.5. Maintenance of Coastal Wetlands, Wetlands Buffers, and Wildlife Corridors	157	
8.6. Protecting Vulnerable Species of Wildlife on Barrier Beaches		
8.7. Aquaculture	158	
8.8. Regional and National Issues that Affect Plum Island Sound	158	

TABLE OF TABLES

Table 2.1. Area of submerged contours for the Parker River-Plum Island Sound Estua	ary,
1965	7
Table 2.2. A summary of the significant morphometric measurements of the area	8
Table 2.3. Soil types in the Plum Island Sound Estuary	11
Table 3.1a. Sampling conditions for the analytes measured during the water quality	
surveys	17
Table 3.1b. Analytical conditions for the analytes measured during the water quality	
surveys	18
Table 3.2. Water Analysis Data Collected at the Little Neck Shore Station (S1), 1965	
Table 3.3. Water Analysis Data Collected at the Bluffs Shore Station (S2), 1965	25
Table 3.4. Water Analysis Data Collected at the Knobs Shore Station (S3), 1965	25
Table 3.5. Water Analysis Data Collected at the Nelson's Island Shore Station (S4), 1	
Table 3.6. Water Analysis Data Collected at the Sub-Headquarters Shore Station (S5	26
1965), 26
Table 3.7. Water Analysis Data Collected at the Town Landing Shore Station (S3), 1	
Tuoie 3.7. Water Thiaryon Bata Conceived at the Town Editating Shore Station (83), T	26
Table 3.8. Ranges for temperature and water analysis measurements for the shoreline	
stations in the Parker River-Plum Island Sound Estuary, 1965	27
Table 3.9. Water Analysis Data Collected at the Camp Sea Haven Offshore Station	27
Table 3.10. Water Analysis Data Collected at the Castle Neck Offshore Station (OS2)	
1965	´´ 27
Table 3.11. Water Analysis Data Collected at the Middle Ground Offshore Station (C	
1965	27
Table 3.12. Water Analysis Data Collected at the Eight Shrimp Trawl Stations (OS4-	
OS11), 1965	28
Table 3.13. Salinity, temperature, and dissolved oxygen (DO) data at the mouth of th	e
Sound off Steep Hill (MAS1), 1992/3	29
Table 3.14. Salinity, temperature, and dissolved oxygen (DO) data at the Ipswich Riv	er
station at Little Neck (MAS 2), 1992/3	30
Table 3.15. Salinity, temperature, and dissolved oxygen (DO) data in the center of Pl	um
Island Sound near Can 23 (MAS12), 1992/3	31
Table 3.16. Salinity, temperature, and dissolved oxygen (DO) data in the Parker Rive	r
near the Newbury Town Landing (MAS15), 1992/3	32
Table 3.17. Salinity, temperature, and dissolved oxygen (DO) data in the Parker Rive	
Route 1 (MAS21), 1992/93	33
Table 3.18. Ranges for temperature, salinity, and dissolved oxygen for the Parker Riv	
Plum Island Sound Estuary, 1992/3	33
Table 3.19. Temperature, salinity, and dissolved oxygen (DO) concentrations from the	ie
mouth of Plum Island Sound in 1989 from a Massachusetts Department of	~ .
environmental Protection survey (station IP10)	34
Table 3.20. Temperature, salinity, and dissolved oxygen from the Ipswich River at Li	
Neck in 1989 (DEP's IP07 station)	34

Table 3.21.	Temperature,	salinity, ar	nd dissolved	oxygen from	Plum
Island Sour	nd off Great N	leck in 198	9 (DEP's IP	09 station)	
	34		•	,	

Table 3.22. Nutrient concentrations from the Steep Hill sampling station (MAS1), 199	
Table 3.23. Nutrient concentrations at the Ipswich River station (MAS2 in 1992, MAS in 1993), 1992/3	36 S3 37
Table 3.24. Nutrient concentrations at Rowley River sampling station (MAS9), 1992/2 Table 3.25. Nutrient concentrations in the middle of Plum Island Sound near Can 23	3 37
(MAS12), 1992/3	38
Table 3.26. Nutrient concentrations at the Parker River by the Newbury Old Town Landing (MAS15), 1992/3	38
Table 3.27. Nutrient concentrations at the Parker River at Route 1 (MAS21), 1992	39
Table 3.28. Nutrient concentrations at the Eagle Hill River (MAS22), 1002 Table 3.29. Nutrient concentrations in the Plum Island River at Jericho Creek (MAS2 1992	39 3), 39
Table 3.30. Chlorophyll α concentrations (mg/L) in selected stations sampled by DEP 1989	o in 40
Table 3.31. Calculated areas and volumes for the model boxes for the 1992 and 1993 surveys	44
Table 3.32. Average salinity by box for the 1992 surveys, with averages by box for surveys 1, 2, 4, and 5	45
Table 3.33. Average salinity by box for the 1993 surveys, with averages for surveys 2	. −5 46
Table 3.34. Calculations for freshwater flow to model box subareas during the 1992 surveys	47
Table 3.35. Calculations for freshwater flow to model box subareas during the 1993 surveys	48
Table 3.36. Model calculated flushing time in days for the two survey years with flush times for the average conditions calculated for four of the surveys in each year	_
Table 3.37. Geometric mean fecal coliform concentrations (as <i>E. coli</i>) in colony formiunits per 100 ml. Plum Island Sound Stations	
Table 3.38. Geometric mean fecal coliform concentrations (as <i>E. coli</i>) in colony form units per 100 ml. Ipswich River and its tributaries	ing 54
Table 3.39. Geometric mean fecal coliform concentrations in the Rowley River and it tributaries. Results are expressed as cfu/100 ml	s 55
Table 3.40. Geometric mean fecal coliform concentrations in cfu/ 100 ml. Parker Riv and its tributaries	er 56
Table 3.41. Geometric mean fecal coliform concentrations in cfu/ 100 ml. Parker Riv NWR	
Table 3.42. Sediment Quality Data (mg/kg dry weight) from the Parker River Watersh Team, Mass (EOEA 1996)	
Table 4.1. Shoreline stations sampled by beach seine Table 4.2. Trawl stations	78 81

Table 4.3. A check list of finfish species collected at all sampling stations in the Parket	er
River-Plum Island Sound Estuary, 1965 (DMF study) and 1993/4 (MAS/WH	
study)	84
Table 4.4. Difference in the catch per unit effort of fish caught by DMF in 1965 and t	he
MAS/Wood Hole study from June 1993 through May 1994	86
Table 4.5. Catch per unit effort (CPUE) of fish using beach seines at Little Neck,	
Ipswich. DMF Station S1. MAS/WH Station 1.	89
Table 4.6. CPUE at Bluffs, Ipswich. DMF Station S2. MAS/WH Station 2	89
Table 4.7. CPUE at Knobs, Rowley. DMF Station S3. MAS/WH Station 3	90
Table 4.8. CPUE at Nelson's Island Shoreline Station. DMF Station S4. MAS/WH	
Station 4	90
Table 4.9. CPUE at Subheadquarters. DMF Station S5. MAS/WH Station 5.	91
Table 4.10. Newbury Town Landing. DMF Station S6. MAS/WH Station 6	91
Table 4.11. Species caught in shrimp trawl by DMF in 1965 and the MAS/WH 1993/	
study of Plum Island Sound	93
Table 4.12. Castle Neck Offshore Station (DMF OS2, MAS/WH Station 10). Total	
number of fish caught over a one year time period	94
Table 4.13. Middle Ground Offshore Station (DMF OS4, MAS/WH Station 12). Total	al
number of fish caught over a one year time period	94
Table 4.14. Frequency of occurrence of macroinvertebrates in seine and trawl sample	
Table 4.15. Dominant organisms associated with different habitats within Plum Island	
Sound	96
Table 4.16. Description and Condition of Fishways in the Parker River, 1997	104
Table 4.17. Boat launchings at the Ipswich Bay Yacht Club in 1965	113
Table 5.1. Commercial licenses issued for shellfishing (from town records)	120
Table 5.2. Recreational licenses issued for shellfishing	120
Table 5.3. Classifications of shellfish beds in Massachusetts	122
Table 5.4. Classifications for Shellfish Areas in Plum Island Sound, Jan 1, 1996	123
Table 5.5. Ipswich landings of soft shell clams in bushels	124
Table 5.6. Bushels of soft shell clams landed in Rowley, 1985-1994	129
Table 5.7. Newbury landings of soft shell clams in bushels	131
Table 5.8. Oyster stocking records	133
Table 6.1. Average of the three highest numbers of birds observed at one time at Plun	
Island during the indicated decade	138
Table 7.1. Major habitats and dominant vegetation on Plum Island	143
Table 8.1. Tidal crossings by town in the Plum Island Sound region	152

TABLE OF FIGURES

Fig. 1.1. Study Area	4
Fig. 2.1. Watersheds of the Parker River/ Plum Island Sound estuary, including the	
Ipswich River	6
Fig. 3.1a. DMF Water quality stations	14
Fig. 3.1b. MAS Water quality stations	15
Fig. 3.2. Location of USGS stream flow gauging stations on the Ipswich and Parker	
Rivers	20
Fig. 3.3. Streamflow in cubic feet per second reported by the Ipswich and Parker USGS	S
gauges for two weeks preceding each of the ten receiving water surveys of Plun	
Island Sound carried out by ASA, Inc.	42
Fig. 3.4. Comparison of mTEC vs MPN methods for fecal coliforms in Plum Island	
Sound	52
Fig. 3.5. Gradient of fecal coliforms – Ipswich River segment	54
Fig. 3.6. Gradient of fecal coliforms – Rowley River segment	55
Fig. 3.7. Gradient of fecal coliforms – Parker River segment	57
Fig. 3.8. Drainage Basin Areas of Plum Island Sound	58
Fig. 3.9. Relative loadings of fecal coliforms to Plum Island Sound from all basins	
59	
Fig. 3.10. Relative loadings of fecal coliforms to Plum Island Sound, Parker River basis	n
only	60
Fig. 3.11. Influence of three upstream stations on fecal coliforms in the lower Parker	
River at station 15. Low tide samples only.	
Fig. 3.11a. Little River at Station 25	61
Fig. 3.11b. Mill River at Station 24	61
Fig. 3.11c. Upstream Parker River at Station 21	62
Fig. 3.12. Influence of Ipswich River on fecal coliform concentrations at the mouth of	
Plum Island Sound. Low tide samples only.	63
Fig. 3.13. Fecal coliforms vs tidal range. Low tide samples.	
Fig. 3.13a. Parker River at Rte 1 (Station 21)	
64	
Fig. 3.13b. Parker River at Rte 1A (Station 15)	65
Fig. 3.13c. Lower Little River station	65
Fig. 3.13d. Lower Mill River station	66
Fig. 3.14. Location of landfills in the Plum Island Sound region	67
Fig. 4.1a. DMF Fish Sampling Stations	79
Fig. 4.1b. MAS/WH Fish Sampling Stations	79
Fig. 4.2a. Distribution of fish caught by seine, DMF 1965 survey	87
Fig. 4.2b. Distribution of fish caught by MAS-WH at shoreline stations 1993/4	87
Fig. 4.3. Yearly comparison of fish numbers over months of both studies	88
Fig. 4.4. Comparison of ecological eveness – Simpson Index	92

Fig. 4.5. Water temperature over time of two studies 97

Fig. 4.6. Salinities over time of two studies	98
Fig. 4.7. Daily discharge at Ipswich and Parker Rivers over time of two studies	99
Fig. 4.8. Location of fishways in the Plum Island Sound area	105
Fig. 4.9. Alewife counts on the Parker River at the Central Street dam carried out by	the /
Parker River Clean Water Association	107
Fig. 4.10. Commercial Striped Bass Landings from Statistical Area #1 (Ipswich Bay	to to
NH border)	110
Fig. 4.11. Public Access Sites	112
Fig. 5.1. Marine Resources – Town of Ipswich (Ipswich River)	125
Fig. 5.2. Marine Resources – Town of Ipswich (Plum Island Sound)	126
Fig. 5.3. Marine Resources – Town of Rowley	128
Fig. 5.4. Marine Resources – Town of Newbury	130
Fig. 8.1. Open Space	149
Fig. 8.2. Tidal Restrictions of Plum Island Sound region	153